

Literature on specific BIVA

Bonaccorsi G, Santomauro F, Lorini C, Indiani L, Pellegrino E, Pasquini G, et al. Mugello Study Working Group. 2016. [Risk of malnutrition in a sample of nonagenarians: Specific versus classic bioelectrical impedance vector analysis](#). *Nutrition*; 32:368-374.

Buffa R, Mereu E, Comandini O, Ibanez ME, Marini E. 2014. [Bioelectrical impedance vectoranalysis \(BIVA\) for the assessment of twocompartment body composition](#). *Eur J Clin Nutr*; 68:1234-1240.

Buffa R, Mereu E, Marini E. 2014. [Reply to the comment on “Lower lean mass and higher percent fat mass in patients with Alzheimer's disease”](#). *Exp Gerontol*; 60:166.

Buffa R, Mereu E, Putzu P, Mereu RM, Marini E. 2014. [Lower lean mass and higher percent fat mass in patients with Alzheimer's disease](#). *Exp Gerontol*; 58:30-33.

Buffa R, Saragat B, Cabras S, Rinaldi AC, Marini E. 2013. [Accuracy of specific BIVA for the assessment of body composition in the United States population](#). *PLoS One*; 8(3):e58533.

Buffa, R., Mereu, E., Succa, V., Latini, V., & Marini, E. 2017. [Specific BIVA recognizes variation of body mass and body composition: Two related but different facets of nutritional status](#). *Nutrition*; 35:1-5.

Camina Martín MA, de Mateo Silleras B, Barrera Ortega S, Domínguez Rodríguez L, Redondo del Río MP. 2014. [Specific bioelectrical impedance vector analysis \(BIVA\) is more accurate than classic BIVA to detect changes in body composition and in nutritional status in institutionalised elderly with dementia](#). *Exp Gerontol*; 57:264-271.

Campa F, Silva AM, Toselli S. 2018. [Changes in Phase Angle and Handgrip Strength Induced by Suspension Training in Older Women](#). *Int J Sports Med*; 39(6):442-449.

Castizo-Olier J, Irurtia A, Jemni M, Carrasco-Marginet M, Fernández-García R, Rodríguez FA. 2018. [Bioelectrical impedance vector analysis \(BIVA\) in sport and exercise: Systematic review and future perspectives](#). *PLoS One*; 7;13(6):e0197957.

Cintosun U, Safer U, Tasci I. 2014. [Comment on “Lower lean mass and higher percent fat mass in patients with Alzheimer's disease”](#). *Exp Gerontol*; 58:303.

Fukuda, D. H., Stout, J. R., Moon, J. R., Smith-Ryan, A. E., Kendall, K. L., & Hoffman, J. R. 2016. [Effects of resistance training on classic and specific bioelectrical impedance vector analysis in elderly women](#). *Exp Gerontol*; 74:9-12.

Ibáñez ME, Mereu E, Buffa R, Gualdi-Russo E, Zaccagni L, Cossu S, et al. 2015. [New specific bioelectrical impedance vector reference values for assessing body composition in the Italian-Spanish young adult population](#). *Am J Hum Biol*.; 27:871-876.

Marini E, Buffa R, Saragat B, Coin A, Toffanello ED, Berton L, Manzato E, Sergi G. 2012. [The potential of classic and specific bioelectrical impedance vector analysis for the assessment of sarcopenia and sarcopenic obesity](#). *Clin Interv Aging*; 7:585-591.

Marini E, Sergi G, Succa V, Saragat B, Sarti S, Coin A, Manzato E, Buffa R. 2013. [Efficacy of specific bioelectrical impedance vector analysis \(BIVA\) for assessing body composition in the elderly.](#) J Nutr Health Aging; 17(6):515-521.

Mereu E, Buffa R, Lussu P, Marini E. 2016. [Phase angle, vector length, and body composition.](#) Am J Clin Nutr; 104:845-847.

Mereu E, Succa V, Buffa R, Sanna C, Mereu RM, Catta O, Marini E. 2018. [Total body and arm bioimpedance in patients with Alzheimer's disease.](#) Exp Gerontol; 102:145-148.

Saragat B, Buffa R, Mereu E, De Rui M, Coin A, Sergi G, Marini E. 2014. [Specific bioelectrical impedance vector reference values for assessing body composition in the Italian elderly.](#) Exp Gerontol; 50:52-56.